

GENERAL MICROBIOLOGY

- Vitamin Requirements of Three Spiroplasmas. C. J. Chang 488-490

STRUCTURE AND FUNCTION

- Protein on the Cell Surface of the Moderately Halophilic Phototrophic Bacterium *Rhodospirillum salexigens*. Dierk Evers, Jürgen Weckesser, and Gerhart Drews 107-111
- Periplasmic Gel: New Concept Resulting from the Reinvestigation of Bacterial Cell Envelope Ultrastructure by New Methods. Jan A. Hobot, Eric Carlemalm, Werner Villiger, and Edward Kellenberger 143-152
- Legionella pneumophila* Surface Antigens Cloned and Expressed in *Escherichia coli* Are Translocated to the Host Cell Surface and Interact with Specific Anti-*Legionella* Antibodies. N. Cary Engleberg, Eric Pearlman, and Barry I. Eisenstein 199-203
- Kinetic Analysis of the Synthesis and Assembly of Type 1 Fimbriae of *Escherichia coli*. Douglas C. Dodd and Barry I. Eisenstein 227-232
- Cell Wall Receptor for Bacteriophage Mu G(+). Rodica Sandulache, Peter Prehm, and Dietmar Kamp 299-303
- Division of Temperature-Sensitive *Streptococcus faecium* Mutants After Return to the Permissive Temperature. Pietro Canepari, Maria Del Mar Lléo, Roberta Fontana, Giuseppe Satta, Gerald D. Shockman, and Lolita Daneo-Moore 427-429
- Isolation of Pigmented and Nonpigmented Mutants of *Serratia marcescens* with Reduced Cell Surface Hydrophobicity. Mel Rosenberg 480-482

PLANT MICROBIOLOGY

- Comparison of T-DNA Oncogene Complements of *Agrobacterium tumefaciens* Tumor-Inducing Plasmids with Limited and Wide Host Ranges. Wallace G. Buchholz and Michael F. Thomashow 319-326
- Host Range Encoded by the *Agrobacterium tumefaciens* Tumor-Inducing Plasmid pTiAg63 Can Be Expanded by Modification of Its T-DNA Oncogene Complement. Wallace G. Buchholz and Michael F. Thomashow 327-332
- Coordinate Expression of Hydrogenase and Ribulose Bisphosphate Carboxylase in *Rhizobium japonicum* Hup^c Mutants. David Merberg and Robert J. Maier 448-450
- Monoclonal Antibodies to *Rhizobium meliloti* and Surface Mutants Insensitive to Them. Eric Johansen, Turlough M. Finan, Malcolm L. Geffer, and Ethan R. Signer 454-457

GENETICS AND MOLECULAR BIOLOGY

- Chloramphenicol-Inducible Gene Expression in *Bacillus subtilis* Is Independent of the Chloramphenicol Acetyltransferase Structural Gene and Its Promoter. Skorn Mongkolsuk, Nicholas P. Ambulos, Jr., and Paul S. Lovett ... 1-8
- Molecular Basis of Isozyme Formation of β -Galactosidases in *Bacillus stearothermophilus*: Isolation of Two β -Galactosidase Genes, *bgaA* and *bgaB*. Haruhisa Hirata, Seiji Negoro, and Hirotsuke Okada 9-14

No. 1	Cloning of the Neutral Protease Gene of <i>Bacillus subtilis</i> and the Use of the Cloned Gene to Create an In Vitro-Derived Deletion Mutation. Maria Y. Yang, Eugenio Ferrari, and Dennis J. Henner	15-21
	Overexpression and Sequence of the <i>Escherichia coli cheY</i> Gene and Biochemical Activities of the CheY Protein. Philip Matsumura, James J. Rydel, Rose Linzmeier, and Dominick Vacante.....	36-41
490	Purification and Characterization of D-Serine Deaminase Activator Protein. Maria C. Heincz, Susan M. Bornstein, and Elizabeth McFall.....	42-49
	Identification and Characterization of a Gene Product That Regulates Type 1 Piliation in <i>Escherichia coli</i> . Paul E. Orndorff and Stanley Falkow	61-66
111	Measurement of In Vivo Expression of the <i>recA</i> Gene of <i>Escherichia coli</i> by Using <i>lacZ</i> Gene Fusions. Jane M. Weisemann, Cindee Funk, and George M. Weinstock	112-121
152	Genetic Characterization and Molecular Cloning of the Tripeptide Permease (<i>tpp</i>) Genes of <i>Salmonella typhimurium</i> . Margaret M. Gibson, Melanie Price, and Christopher F. Higgins	122-130
203	Anaerobic and Leucine-Dependent Expression of a Peptide Transport Gene in <i>Salmonella typhimurium</i> . Derek J. Jamieson and Christopher F. Higgins	131-136
232	Cloning and Characterization of <i>recA</i> Genes from <i>Proteus vulgaris</i> , <i>Erwinia carotovora</i> , <i>Shigella flexneri</i> , and <i>Escherichia coli</i> B/r. Sherry L. Keener, Kevin P. McNamee and Kevin McEntee	153-160
303	Study of Regulation and Transport of Hemolysin by Using Fusion of the β -Galactosidase Gene (<i>lacZ</i>) to Hemolysin Genes. Antonio Juarez, Michael Härtlein, and Werner Goebel	161-168
429	Cloning of the <i>tyrP</i> Gene and Further Characterization of the Tyrosine-Specific Transport System in <i>Escherichia coli</i> K-12. Peter J. Wookey, James Pittard, Susan M. Forrest, and Barrie E. Davidson.....	169-174
482	Construction of a <i>tyrP-lac</i> Operon Fusion Strain and Its Use in the Isolation and Analysis of Mutants Derepressed for <i>tyrP</i> Expression. Paul A. Kasian and James Pittard	175-183
	<i>lon</i> Transcriptional Regulation of Genes Necessary for Capsular Polysaccharide Synthesis in <i>Escherichia coli</i> K-12. Patsy Trisler and Susan Gottesman....	184-191
326	Regulation of the F-Factor <i>pif</i> Operon: <i>pifO</i> , a Site Required in <i>cis</i> for Autoregulation, Titrates the <i>pifC</i> Product in <i>trans</i> . Jeff F. Miller and Michael H. Malamy	192-198
332	Modification of UV-Induced Mutation Frequency and Cell Survival of <i>Escherichia coli</i> B/r WP2 <i>trpE65</i> by Treatment Before Irradiation. Charles O. Doudney and Charles N. Rinaldi	233-238
450	Molecular Cloning of the Hemolysin Determinant from <i>Vibrio cholerae</i> El Tor. Steven L. Goldberg and John R. Murphy	239-244
457	Genetic and Physical Analyses of <i>Caulobacter crescentus trp</i> Genes. Malcolm E. Winkler, Patricia V. Schoenlein, Carolyn M. Ross, John T. Barrett, and Bert Ely	279-287
	Mutagenesis of <i>Neisseria gonorrhoeae</i> : Absence of Error-Prone Repair. L. A. Campbell and R. E. Yasbin.....	288-293
1-8	Genetic System for Analyzing <i>Escherichia coli</i> Thymidylate Synthase. Marlene Belfort and Joan Pedersen-Lane.....	371-378
-14	Identification and Regulation of the <i>glnL</i> Operator-Promoter of the Complex <i>glnALG</i> Operon of <i>Escherichia coli</i> . Shizue Ueno-Nishio, Susan Mango, Lawrence J. Reitzer, and Boris Magasanik	379-384

Enzymatic and Nucleotide Sequence Studies of a Kanamycin-Inactivating Enzyme Encoded by a Plasmid from Thermophilic Bacilli in Comparison with That Encoded by Plasmid pUB110. Masazumi Matsumura, Yoshio Katakura, Tadayuki Imanaka, and Shuichi Aiba	413-420
Construction of a <i>Bacillus subtilis</i> Double Mutant Deficient in Extracellular Alkaline and Neutral Proteases. Fujio Kawamura and Roy H. Doi.....	442-444
Characterization of Interspecific Plasmid Transfer Mediated by <i>Bacillus subtilis</i> Temperate Bacteriophage SP02. Robert Marrero, Frank E. Young, and Ronald E. Yasbin.....	458-461

PLASMIDS AND TRANSPOSONS

IncFII Plasmid Incompatibility Product and Its Target Are Both RNA Transcripts. David D. Womble, Xinnian Dong, Ru Ping Wu, Verne A. Luckow, Asuncion F. Martinez, and Robert H. Rownd.....	28-35
Transposon Tn10-Like Tetracycline Resistance Determinants in <i>Haemophilus parainfluenzae</i>. Stuart B. Levy, Annie Buu-Hoi, and Bonnie Marshall	87-94
Inverted Repeat Sequences Flank a <i>Bacillus thuringiensis</i> Crystal Protein Gene. James W. Kronstad and H. R. Whiteley	95-102
Second Replicon in ColV2-K94 Mediates the Stable Coexistence of Two Incompatible Plasmids. Peter C. Weber, Gopa Mitra, and Sunil Palchaudhuri.....	245-250
Transposon Mutagenesis Analysis of <i>meta</i>-Cleavage Pathway Operon Genes of the TOL Plasmid of <i>Pseudomonas putida</i> mt-2. Shigeaki Harayama, Philip R. Lehrbach, and Kenneth N. Timmis.....	251-255
Aerobactin Iron Uptake Sequences in Plasmid ColV-K30 Are Flanked by Inverted IS1-Like Elements and Replication Regions. Jose F. Perez-Casal and Jorge H. Crosa	256-265
Aerobactin Genes in <i>Shigella</i> spp. Kathleen M. Lawlor and Shelley M. Payne	266-272
DNA Sequence of the F <i>tra</i>LE Region That Includes the Gene for F Pilin. Laura S. Frost, W. Paranchych, and Neil S. Willetts	395-401
Localization, Cloning, and Sequence Determination of the Conjugative Plasmid ColB2 Pilin Gene. B. Brett Finlay, Laura S. Frost, and William Paranchych	402-407
Polypeptide Involved in the <i>Escherichia coli</i> Plasmid-Mediated Citrate Transport System. Toru Hirato, Morikazu Shinagawa, Naotaka Ishiguro, and Gihei Sato.....	421-426
High Incidence of Transposon Tn3 Insertions into a Replication Control Gene of the Chimeric R/Ent Plasmid pCG86 of <i>Escherichia coli</i>. Roger N. Picken, Anthony J. Mazaitis, and Werner K. Maas	430-433
In Vitro Deletions in the Partition Locus of Plasmid pSC101. S. Roger Kalla and Petter Gustafsson	434-437
Conjugal Transfer of Bacterial Chromosomes Mediated by the RK2 Plasmid Transfer Origin Cloned into Transposon Tn5. Emanuel A. Jakobson and Donald G. Guiney, Jr.	451-453
Mobilization of a Sym Plasmid from a Fast-Growing Cowpea <i>Rhizobium</i> Strain. Nigel A. Morrison, Ying Hua Cen, Hen Cai Chen, Jacek Plazinski, Robert Ridge, and Barry G. Rolfe	483-487

EUCARYOTIC CELLS

Effect of Castanospermine on the Structure and Secretion of Glycoprotein Enzymes in <i>Aspergillus fumigatus</i>. Alan D. Elbein, Michael Mitchell, and Russell J. Molyneux	67-75
--	-------

	Secretion Can Proceed Uncoupled from Net Plasma Membrane Expansion in Inositol-Starved <i>Saccharomyces cerevisiae</i> . Katharine D. Atkinson and Robert M. Ramirez	80-86
420	<i>Saccharomyces cerevisiae</i> Aldolase Mutants. Zita Lobo	222-226
444	Possible Relationship Between Cyclic AMP and Idiophasic Metabolism in the White Rot Fungus <i>Phanerochaete chrysosporium</i> . Malcolm J. MacDonald, Alistair Paterson, and Paul Broda	470-472

MEMBRANES

461	Proline Uptake Through the Major Transport System of <i>Salmonella typhimurium</i> Is Coupled to Sodium Ions. John Cairney, Christopher F. Higgins, and Ian R. Booth	22-27
-35	Methylammonium Transport in <i>Anacystis nidulans</i> R-2. Samy Boussiba, Waltraud Dilling, and Jane Gibson	204-210
-94	<i>Escherichia coli</i> Mutants with Altered Cation Recognition by the Melibiose Carrier. Sumiko Shiota, Hiromichi Yazyu, and Tomofusa Tsuchiya	445-447
102	Respiratory System of Vegetative and Sporulating <i>Bacillus cereus</i> . Jose E. Escamilla and Maria C. Benito	473-477

PHYSIOLOGY AND METABOLISM

250	Levels of Cyclic-2,3-Diphosphoglycerate in <i>Methanobacterium thermoautotrophicum</i> During Phosphate Limitation. Robert J. Seely and David E. Fahrney	50-54
255	Regulatory role of Phosphate and Other Anions in Transport of ADP and ATP by <i>Rickettsia prowazekii</i> . Herbert H. Winkler and Robin M. Daugherty	76-79
265	Hydrogenase Activity in <i>Azospirillum brasilense</i> Is Inhibited by Nitrite, Nitric Oxide, Carbon Monoxide, and Acetylene. Karl H. Tibelius and Roger Knowles	103-106
272	Induction of Superoxide Dismutases in <i>Escherichia coli</i> by Manganese and Iron. Shirley Y. R. Pugh, James L. DiGuseppi, and Irwin Fridovich	137-142
401	Futile Xylitol Cycle in <i>Lactobacillus casei</i> . Sally Z. Hausman, John Thompson, and Jack London	211-215
407	H ⁺ /ATP Stoichiometry of Cowpea <i>Rhizobium</i> sp. Strain 32H1 Cells Grown Under Nitrogen-Fixing and Nitrogen-Nonfixing Conditions. J. W. Gober and E. R. Kashket	216-221
426	Transport of Mevalonate by <i>Pseudomonas</i> sp. Strain M. John F. Gill, Jr., Michael J. Beach, and Victor W. Rodwell	294-298
433	Specificity and Mechanism of Ferrioxamine-Mediated Iron Transport in <i>Streptomyces pilosus</i> . Gertraud Müller and Kenneth N. Raymond	304-312
437	Iron Transport in <i>Streptomyces pilosus</i> Mediated by Ferrichrome Siderophores, Rhodotorulic Acid, and Enantio-Rhodotorulic Acid. Gertraud Müller, Berthold F. Matzanke, and Kenneth N. Raymond	313-318
453	Properties of ATP-Dependent Protein Kinase from <i>Streptococcus pyogenes</i> That Phosphorylates a Seryl Residue in HPr, a Phosphocarrier Protein of the Phosphotransferase System. Jonathan Reizer, Michael J. Novotny, Wolfgang Hengstenberg, and Milton H. Saier, Jr.	333-340
487	Permeability Changes in the Cytoplasmic Membrane of <i>Escherichia coli</i> K-12 Early After Infection with Bacteriophage T1. Heribert Keweloh and Evert P. Bakker	347-353
-75	Increased Permeability and Subsequent Resealing of the Host Cell Membrane Early After Infection of <i>Escherichia coli</i> with Bacteriophage T1. Heribert W. Keweloh and Evert P. Bakker	354-359

Role of III^{Glc} of the Phosphoenolpyruvate-Glucose Phosphotransferase System in Inducer Exclusion in <i>Escherichia coli</i>. Stephen O. Nelson, Joseph Lengeler, and Pieter W. Postma	360-364
Methanogenic Cleavage of Acetate by Lysates of <i>Methanosarcina barkeri</i>. Larry Baresi	365-370
Effect of Chloride and Glutamate Ions on In Vitro Protein Synthesis by the Moderate Halophile <i>Vibrio costicola</i>. Masahiro Kamekura and Donn J. Kushner	385-390
<i>ilvB</i>-Encoded Acetolactate Synthase Is Resistant to the Herbicide Sulfometuron Methyl. Robert A. LaRossa and Dana R. Smulski	391-394
Novel Inducers of the Xylan-Degrading Enzyme System of <i>Cryptococcus albidus</i>. Peter Biely and Eva Petráková	408-412
Single, Chemically Defined Sporulation Medium for <i>Bacillus subtilis</i>: Growth, Sporulation, and Extracellular Protease Production. James H. Hageman, Gary W. Shankweiler, Pamela R. Wall, Karen Franich, Gloria W. McCowan, Susan M. Cauble, Jesus Grajeda, and Casilda Quinones	438-441
Physiological Role of Pyruvate Oxidase in the Aerobic Metabolism of <i>Lactobacillus plantarum</i>. Barbara Sedewitz, Karl Heinz Schleifer, and Friedrich Götz	462-465
Effects of Cultivation Gas Phase on Hydrogenase of the Acetogen <i>Clostridium thermoaceticum</i>. Rebecca Kellum and Harold L. Drake	466-469
Betaine Is the Main Compatible Solute of Halophilic Eubacteria. Johannes F. Imhoff and F. Rodriguez-Valera	478-479

ENZYMOLOGY

Purification and Properties of Dihydroxyacetone Kinase from <i>Klebsiella pneumoniae</i>. E. A. Johnson, S. K. Burke, R. G. Forage, and E. C. C. Lin	55-60
Purification and Biochemical Characterization of Pyruvate Oxidase from <i>Lactobacillus plantarum</i>. Barbara Sedewitz, Karl Heinz Schleifer, and Friedrich Götz	273-278
Purification and Properties of γ-Glutamyltranspeptidase from <i>Proteus mirabilis</i>. Reiko Nakayama, Hidehiko Kumagai, and Tatsurokuro Tochikura	341-346

ERRATA

Changes in the Composition of <i>Escherichia coli</i> Murein as It Ages During Exponential Growth. Lars G. Burman and James T. Park	491
Post-transcriptional Regulation of Chloramphenicol Acetyl Transferase. Woo-Hyeon Byeon and Bernard Weisblum	491

Aerobactin Genes in *Shigella* spp.

KATHLEEN M. LAWLOR AND SHELLEY M. PAYNE*

Department of Microbiology, University of Texas, Austin, Texas 78712-1095

Received 18 June 1984/Accepted 28 July 1984

Aerobactin, a hydroxamate iron transport compound, is synthesized by some, but not all, *Shigella* species. Conjugation and hybridization studies indicated that the genes for the synthesis and transport of aerobactin are linked and are found on the chromosome of *Shigella flexneri*, *S. boydii*, and *S. sonnei*. The genes were not found in *S. dysenteriae*. A number of aerobactin synthesis mutants and transport mutants have been isolated. The most common mutations are deletions of the biosynthesis or biosynthesis and transport genes. The *Shigella* aerobactin genes share considerable homology with the *E. coli* ColV aerobactin genes. On the ColV plasmid and in the *Shigella* chromosome, the aerobactin genes are associated with a repetitive sequence which has been identified as ISI.

Shigella species synthesize both phenolate and hydroxamate siderophores for acquisition of iron. *Shigella sonnei* utilizes enterobactin (enterochelin) (25), a phenolate siderophore common to many enteric species (19), and some strains produce an additional hydroxamate siderophore (25). Similarly, *S. boydii* has been reported to produce both phenolate and hydroxamate siderophores (25). *S. flexneri* strains normally synthesize only aerobactin (22), a secondary hydroxamate originally isolated from cultures of *Aerobacter aerogenes* (9) and more recently from *Escherichia coli* ColV strains (29). Rare isolates and laboratory variants of *S. flexneri* produce enterobactin in addition to aerobactin (23).

The genetics of iron transport in enteric bacteria has been studied primarily in *E. coli* and *Salmonella typhimurium*. The genes for enterobactin synthesis and transport are linked and are carried on the chromosome in both species (12-14, 26). Aerobactin genes, however, are encoded by the ColV plasmid in *E. coli* (29) and by a large plasmid in *A. aerogenes* (18). In other species, the location of the aerobactin genes is unclear.

Since differences were observed in the types of iron transport systems expressed by different *Shigella* species (22, 23, 25) and even by members of the same species, it was of interest to examine the genetics of siderophore synthesis and transport in *Shigella* spp. and to compare it to the ColV system.

MATERIALS AND METHODS

Strains and media. Bacterial strains, plasmids, a phage, and their sources are listed in Table 1. All other *Shigella* isolates were provided by Gloria Pierce, Texas Department of Health, Austin, Texas. Plasmid pKLS1 was a spontaneous deletion of pABN1. This plasmid was isolated by selecting ampicillin-resistant, cloacin-resistant mutants of pABN1. Ampicillin selects for maintenance of the pABN1 vector, whereas cloacin selects for loss of the 74K aerobactin receptor protein (3). This plasmid contains 1.7 kilobases (kb) of the original 16.3 kb of ColV sequences as determined by restriction enzyme digestion and hybridization with pABN1. The subclone pKLS10 was constructed by *Bgl*II restriction enzyme digestion and ligation of electroeluted fragments of pABN1. Both plasmids have the original vec-

tor, pPlac, from pABN1. Bacterial stocks were maintained frozen at -70°C in tryptic soy broth (Difco Laboratories) with 20% glycerol. Luria broth (15) was used for growth of *Shigella* strains. Low-iron Tris-buffered medium without added iron was used as described previously (22, 23) to determine siderophore production.

Siderophore assays. Strains were grown through two passages of Tris medium to induce siderophore synthesis. Phenolates were assayed by the colorimetric assay of Arnov (1), and hydroxamates were assayed by the ferric perchlorate assay of Atkin et al. (2) or the more sensitive Csaky test (6). Transport of siderophores was determined by a bioassay (22, 23) in which the ability of a siderophore to stimulate growth of bacteria in low-iron agar medium was determined. Siderophores were isolated by extraction with organic solvents and chromatography and were compared with aerobactin or enterobactin by thin-layer chromatography as previously described (22, 23).

Southern hybridization. Chromosomal DNA was prepared by the procedure of Marmur (17). *Shigella* plasmid DNA was isolated by the method of Hansen and Olsen (10), a procedure which allows isolation of large plasmids. Cleared lysates of sodium dodecyl sulfate-lysed cells (15) were the source of other plasmids. DNA was cut with restriction endonucleases (New England Biolabs) and electrophoresed through 0.9% agarose gels. DNA fragments were transferred to nitrocellulose (Schleicher & Schuell, Inc.) by the method of Southern (28) and were hybridized by that of Maniatis et al. (15). Hybridization probes were prepared by restriction enzyme digestion of plasmid or phage DNA and separation of insert fragments from vector on agarose gels. It was necessary to eliminate vector sequences since some of the small *Shigella* plasmids which frequently contaminate chromosomal DNA preparations were found to have homology with ColE1 vectors. DNA fragments were electroeluted into wells in the agarose filled with 50% glycerol in electrophoresis buffer. Eluted fragments were labeled with [³²P]dCTP (New England Nuclear Corp.) by nick translation (16). Only labeled fragments which did not show significant hybridization to vector DNA were used. Filters were hybridized at 68°C and washed at 68°C with 0.1 × SSC (SSC is 0.15 M NaCl plus 0.015 M sodium citrate) for stringent hybridization conditions (15). The SSC concentration was increased to 1× when less stringent conditions were desired.

Conjugation. Overnight cultures of the donor (*S. flexneri* 256) and recipient (SA255) were diluted in Luria broth and

* Corresponding author.

TABLE 1. Bacterial strains, plasmids, and phage

Strain ^a	Source/Reference
Bacteria	
<i>S. flexneri</i>	
SA100 Hds ⁺ Hdu ⁺	23
SA201 Hds ⁻	23
SA255 Hds ⁻ Hdu ⁻	23
SA845 (F ⁺ /tet ^r)	This study
256 (Hfr)	R. Neill, Walter Reed Army Institute for Research (8)
<i>S. dysenteriae</i> 4576	Texas Department of Health
<i>S. sonnei</i> 1245	Texas Department of Health
<i>S. boydii</i> 1392	Texas Department of Health
Phage	
VAX3 (IS1)	7
Plasmids	
ColV-K30	J. H. Crosa (29)
pABN1	4
pKLS10	Subclone of pABN1
pKLS1	Deletion plasmid derived from pABN1

^a Hds, Aerobactin synthesis; Hdu, aerobactin utilization.

grown to mid-log phase. Equal volumes (1 ml) were mixed in a 125-ml flask and were incubated at 37°C for 2 h. The mixture was diluted and plated on Luria broth agar containing an antibiotic to select again the donor and 25 µg of deferrated ethylenediamine-di-(*o*-hydroxyphenylacetic acid) (EDDA; Sigma Chemical Co.) per ml (27). The iron chelator EDDA inhibits growth of cells which lack a siderophore-mediated iron transport system.

TABLE 2. Synthesis and utilization of siderophores by *Shigella* species

Species	Aerobactin		Enterobactin	
	Syn-thesis ^a	Utiliza-tion ^b	Syn-thesis ^c	Utiliza-tion ^b
<i>S. boydii</i> 1392	+	(0.08)	+	—
<i>S. dysenteriae</i> 4576	—	—	+	(0.16)
<i>S. flexneri</i> SA100	+	(0.07)	+	—
<i>S. sonnei</i> 1245	—	—	+	(0.27)

^a Aerobactin synthesis determined by ferric perchlorate (2) and Csaky (6) assays and comparison of the isolated siderophore to aerobactin on thin-layer chromatography. Numbers in parentheses show the absorbance at 500 nm (by ferric perchlorate assay) of T medium culture supernatant.

^b Determined by bioassay.

^c Enterobactin synthesis determined by Arnow (1) assay and comparison of isolated phenolate to enterobactin on thin-layer chromatography. Numbers in parentheses show the absorbance at 515 nm (by Arnow assay) of T medium culture supernatant.

RESULTS

Strains of *Shigella* spp. were examined to determine the presence and location of the aerobactin genes. Initially, strains were grown in low-iron medium and tested for the ability to synthesize and utilize aerobactin and enterobactin. Results of representative strains of each species are shown in Table 2. In all, 2 strains of *S. dysenteriae*, 10 strains of *S. sonnei*, and 3 strains of *S. boydii* were tested. One hundred strains of *S. flexneri* had been assayed previously (23), and four additional strains were included in this study. No evidence for the secretion of hydroxamates by *S. dysenteriae* or *S. sonnei* was found. The ferric perchlorate (2) and Csaky (6) assays were negative, and no functional aerobactin was detected by bioassay. Both species were found to secrete and utilize enterobactin. Although the *S. sonnei* strains tested did not synthesize aerobactin, they were able to transport the siderophore and utilize it for growth in low-




FIG. 1. Partial restriction map of ColV-K30 and hybridization probes. Restriction endonucleases: B, *Bam*HI; Bg, *Bgl*II; E, *Eco*RI; H, *Hind*III; P, *Pst*I; S, *Sal*I. *Pst*I and *Bgl*II sites are incomplete. (A) ColV-K30. Solid bar indicates the regions in which IS1 sequences map. Locations of aerobactin synthesis genes and 74K aerobactin receptor are indicated (4, 18). (B) The 16.3-kb *Hind*III fragment of ColV DNA in pABN1. This insert contains both aerobactin biosynthesis and transport genes (4). (C) *Hind*III-*Bgl*II subclone of pABN1 containing aerobactin biosynthesis genes. (D) *Hind*III fragment of pKLS1, a deletion mutant of pABN1. Dotted line indicates deleted sequences. (E) A 6.4-kb *Bam*HI fragment of pABN1 containing one copy of IS1.

iron medium, indicating expression of the gene(s) for the aerobactin receptor. The strains of *S. boydii* tested were found to synthesize and transport aerobactin. Neither synthesis nor utilization of enterobactin occurred in these strains.

To determine the presence of aerobactin genes directly, chromosomal and plasmid DNA isolated from strains of all four species was hybridized with the aerobactin genes cloned from the *E. coli* ColV plasmid (4). The 16.3-kb *Hind*III fragment of plasmid pABN1, which contains the aerobactin biosynthesis and transport genes (4) (Fig. 1B), was labeled by nick translation and hybridized to Southern blots of restriction enzyme-digested *Shigella* DNA. A large number of DNA fragments of *S. dysenteriae*, *S. sonnei*, and *S. flexneri* DNA, ranging in size from 0.9 to 23 kb, hybridized to the ColV sequences (Fig. 2, lanes B through D). The *S. flexneri* fragments could be seen more clearly when the filter was exposed for a longer period of time (data not shown), which overexposed the *S. dysenteriae* and *S. sonnei* fragments. The difference in intensity of *S. flexneri* fragments and those of *S. dysenteriae* and *S. sonnei* may reflect differences in copy number or in homology. Fewer fragments of *S. boydii* DNA were detected (Fig. 2A). Since the ColV probe was relatively large, it was necessary to use fragments or subclones of pABN1 (Fig. 1C and E) to determine which of the multiple bands represented aerobactin genes and whether the remaining bands represented a repeated sequence. One subclone, pKLS10, contained a 4.8-kb *Hind*III-*Bgl*II fragment of pABN1 (Fig. 1C). Iron transport mutants of *S. flexneri* were transformed with this plasmid to determine whether it contained aerobactin genes. SA201, a mutant defective in synthesis of aerobactin, was complemented by this plasmid, and aerobactin was detected in supernatant fluids of the transformant. SA255, a mutant defective in both synthesis and transport of aerobactin, was not complemented by pKLS10. Hybridization to this plasmid was used to detect aerobactin biosynthesis genes. When the *Hind*III-*Bgl*II fragment of this plasmid was used, a single

*Hind*III (Fig. 2, lanes E, F, and H) or *Eco*RI (data not shown) fragment of *S. flexneri*, *S. sonnei*, or *S. boydii* hybridized under stringent conditions. The *Hind*III fragment was approximately 12.4 kb in *S. flexneri* and *S. sonnei* and approximately 11.0 kb in *S. boydii*. No hybridization to *S. dysenteriae* was detected (Fig. 2, lane G), even when less stringent conditions were used (data not shown).

Since aerobactin genes previously had been found to be plasmid encoded (29), *Shigella* plasmids were also hybridized to ColV aerobactin genes. Cleared lysates (10) of all four species contained one or more plasmids ranging in size from 1.5 to 240 kb, but none hybridized to the *Hind*III-*Bgl*II fragment of pKLS10 (Fig. 3).

The multiple bands detected when the entire *Hind*III fragment of pABN1 was used were also seen when a 6.4-kb *Bam*HI fragment of pABN1 (Fig. 1E) was hybridized to *Shigella* chromosomal DNA (Fig. 2, lanes I through L). This *Bam*HI probe shares 1.0 kb with pKLS10 and includes an additional 5.4 kb as determined by restriction enzyme digestion and hybridization.

The large number of fragments hybridizing to the *Bam*HI probe suggested that an insertion sequence might be present. Since a large number of copies (>40) of IS1 are found in most *Shigella* species (20), it appeared likely that the sequence might be IS1. A probe containing IS1 was used to determine whether this insertion sequence was present on pABN1. VAX3, a λ phage which acquired a copy of IS1 from R100 (7), was used as the source of IS1 DNA. The 10.2-kb *Hind*III fragment of the phage containing IS1 was hybridized to pABN1, and a single copy of IS1 was found on the 6.4-kb *Bam*HI fragment (Fig. 4, lane C). No hybridization between wild-type λ and pABN1 was seen (data not shown), indicating that the homology between pABN1 and VAX3 was due to IS1 and not to the λ sequences. Digestion of pABN1 with *Pst*I produced two fragments which hybridized to IS1 (Fig. 4, lane A), as was expected since there is a single *Pst*I site within IS1 (21).

To determine whether additional copies of IS1 were




FIG. 2. Hybridization of ColV sequences to *Shigella* DNA. DNA of *S. boydii* (lanes A, E, and I), *S. sonnei* (lanes B, F, and J), and *S. dysenteriae* (lanes C, G, and K), and *S. flexneri* (lanes D, H, and L) was cut with *Hind*III and hybridized to the 16.3-kb *Hind*III fragment of pABN1 (lanes A through D), the 4.8-kb *Hind*III-*Bgl*II fragment of pKLS10 (lanes E through H), or the 6.4-kb *Bam*HI fragment of pABN1 (lanes I through L). Positions of molecular size standards are indicated at the left.




FIG. 3. Hybridization of ColV sequences to *Shigella* plasmids. (A) Plasmids of *S. flexneri* (lane 1), *S. sonnei* (lane 2), *S. boydii* (lane 3), and *S. dysenteriae* (lane 4) were electrophoresed in 0.9% agarose and were stained with ethidium bromide. A Southern blot of this gel (B) was hybridized to the 4.8-kb insert of pKLS10; lane numbers are as in (A). A sample of *S. flexneri* chromosomal DNA cut with *Hind*III (C) was included on the gel to ensure that hybridization would be detected. Sizes of *S. flexneri* plasmids are indicated at the left.

associated with the aerobactin genes, restriction enzyme fragments of the entire ColV plasmid were hybridized to pKLS10 (Fig. 5, lanes B and E) and VA λ 3 (Fig. 5, lane G). A single 8.4-kb *Eco*RI fragment of ColV (Fig. 5, lane E) and a single 16.3-kb *Hind*III fragment (data not shown) were detected by using the aerobactin gene probe pKLS10. Two fragments were detected when the plasmid was cut with *Sal*I (Fig. 5, lane B) or *Bam*HI (data not shown), both of which cut within this region. The intensity of hybridization of the two *Sal*I fragments of ColV DNA (Fig. 5B) differs because *Sal*I cuts very near the end of the region detected by pKLS10 (Fig. 1), leaving only a small region of homology between pKLS10 and the 21-kb *Sal*I fragment.

A duplicate Southern blot was hybridized with the IS1 probe to determine whether the ColV DNA fragments which hybridized to the aerobactin gene probe also contained IS1 (Fig. 5). Four of the *Sal*I fragments (Fig. 5, lane G) and three of the *Eco*RI fragments (data not shown) hybridized to the IS1 probe. Since these enzymes do not cut within IS1 (21), there are at least four copies of the insertion sequence on the ColV plasmid. The different intensities of the four bands may indicate more than one copy of IS1 on some fragments, or there may be differences in the extent of homology between ColV insertion sequences and the probe. Two copies of the insertion sequence are on 21- and 12.5-kb *Sal*I fragments, which also hybridize to the aerobactin gene probe, pKLS10 (Fig. 5, lanes B and G). One of these copies of IS1 is contained on pABN1 (Fig. 4) but is absent from pKLS1 (Fig. 5, lane H), a deletion mutant of pABN1 which lacks the 12.5-kb *Sal*I fragment (Fig. 1D). This copy of IS1 has been mapped 3' to the aerobactin biosynthesis genes by enzyme digestion and hybridization (Fig. 4). A second copy lies 5' to the aerobactin genes on the 21-kb *Sal*I fragment. This copy is to the left of the *Eco*RI site at position 20 kb on the ColV map (Fig. 1) since the 8.4-kb *Eco*RI fragment of ColV DNA which hybridized to pKLS10 did not hybridize to IS1 (data not shown).

Hybridization of the ColV aerobactin genes (pKLS10) to *S. flexneri* and *S. boydii* DNA (Fig. 5) indicate considerable homology. *Eco*RI fragments of the same size, 8.4 kb, were detected in all three species under stringent conditions (Fig. 5, lanes D, E, and F). The flanking sequences are different, however, and the sizes of *Sal*I fragments (Fig. 5, lanes A, B,

and C) and *Bam*HI fragments (data not shown) differ in these species.

Hybridization of IS1 to fragments of *S. flexneri* and *S. boydii* (data not shown) produced the same pattern obtained with the 6.4-kb *Bam*HI fragment of pABN1 (Fig. 2, lanes L and I). However, the large number of fragments in the chromosomal digests and the numerous copies of IS1 in *S. flexneri* made it impossible to determine whether fragments of the same size which hybridize to both the IS1 and aerobactin gene probes were in fact the same fragment.

If IS1 is associated with the aerobactin genes in the




FIG. 4. IS1 sequences of pABN1. pABN1 was cut with *Pst*I (lane A), *Bgl*II (lane B), or *Bam*HI (lane C) and hybridized to the 10.2-kb *Hind*III fragment of VA λ 3 containing IS1. Positions of molecular size standards are indicated at the left.




FIG. 5. Hybridization of aerobactin sequences to ColV, *S. flexneri*, and *S. boydii* and hybridization of IS1 to ColV. DNA of *S. flexneri* (lanes A and D), *S. boydii* (lanes C and F), and ColV-K30 (lanes B and E) was cut with *Sall* (lanes A through C) or *EcoRI* (lanes D through F) and hybridized to the 4.8-kb *HindIII*-*BglIII* insert of pKLS10 containing aerobactin sequences. In addition, ColV-K30 (lane G) and pKLS1 (lane H) DNA were cut with *Sall* and hybridized to the 10.2-kb *HindIII* fragment of V λ 3 containing IS1. Arrows indicate fragments which hybridized to both the aerobactin and IS1 probes. Positions of standards, indicated at left, were the same for both gels.

Shigella chromosome as it is with the ColV aerobactin genes, a relatively high frequency of deletion mutations might be expected to occur in this region (5). Thirty-three independent, spontaneous mutants which failed to grow in low-iron medium were isolated from *S. flexneri* SA100 and were tested for synthesis and utilization of aerobactin. Twenty-six of these mutants failed to synthesize detectable aerobactin but were able to transport the compound; six mutants neither synthesized nor transported aerobactin. An additional mutant, SA280, was found to be defective in transport, but not in synthesis of the siderophore. This mutant reverted to wild type at a frequency of approximately 10^{-6} . None of the synthesis or synthesis and transport mutants reverted at a detectable frequency. This suggests that the majority of these mutants are deletions. DNA was isolated from two of the mutants, SA201 (a synthesis mutant) and SA255 (a synthesis and transport mutant), and was hybridized to pKLS10 (Fig. 6). SA201 DNA hybridized to pKLS10, but the size of the *HindIII* fragment was altered and the intensity was diminished compared with parental DNA (Fig. 6, lanes A and B). SA255 has no homology with pKLS10 (Fig. 6, lane C), indicating a deletion of aerobactin sequences. If the deletion of aerobactin genes in SA255 was due to IS1, the deletion should extend to the insertion sequence and alter the size of any restriction fragment containing IS1 and flanking sequences. An IS1 adjacent to the aerobactin sequences would not necessarily be on the same restriction fragment as the aerobactin genes. DNA from SA100 and SA255 was cut with *HindIII*, which does not cut within IS1, and hybridized to V λ 3. Two restriction fragments of 2.5 and 3.8 kb which were present in SA100 were absent from SA255 (Fig. 7). This indicates an alteration in sequences adjacent to the insertion sequence or deletion of the insertion sequence. The gel was exposed for a shorter

period of time to determine changes in hybridization to larger fragments, but no differences were seen. Deletion of aerobactin genes as well as sequences adjacent to two copies of IS1 suggests that the aerobactin genes of *S. flexneri*, like ColV, are flanked by copies of IS1. Deletions extending from one copy of IS1 into the aerobactin genes or recombination between two copies of IS1 would result in mutations of the type found in SA255. Smaller deletions would produce mutants defective in biosynthesis only.

Conjugation between an aerobactin-producing (Hds⁺




FIG. 6. Hybridization of aerobactin genes to *S. flexneri* mutants. DNA of *S. flexneri* SA100 (lane A), SA201 (lane B), and SA255 (lane C) was cut with *HindIII* and hybridized to the insert of pKLS10. Sizes of standards are indicated at the left.




FIG. 7. Hybridization of *IS1* to *S. flexneri*. DNA of SA100 (lane A) and SA255, an aerobactin biosynthesis and transport mutant of SA100 (lane B), was cut with *Hind*III and hybridized to the 10.2-kb *Hind*III fragment of VA λ 3 containing *IS1*. Arrows indicate bands which are absent from SA255 DNA.

Hdu⁺) donor and a streptomycin-resistant derivative of the deletion mutant SA255 was used to confirm the chromosomal location of these genes. The donor, strain 256, transfers the chromosome in a clockwise direction from approximately 7 min by analogy with the *E. coli* map (8). Hds⁺ Hdu⁺ recombinants were obtained at frequencies of 10⁻⁴ to 10⁻⁵. This frequency was higher than that observed for other markers (10⁻⁵ to 10⁻⁸), suggesting that the genes map near the origin of transfer. Linkage to other markers in this area, however, was not consistent. Additional studies will be done to determine the location of these genes.

These experiments were repeated with an F donor, SA845, instead of the Hfr donor. No Hds⁺ Hdu⁺ recombinants were obtained from these matings. Transfer by an Hfr but not an F donor indicates that the genes are chromosomal.

DISCUSSION

S. flexneri utilizes a hydroxamate compound, aerobactin, for transport of iron. This siderophore also has been isolated from other enteric bacteria. Although all strains of *S. flexneri* appear to use this compound, the aerobactin system is not uniformly present in other species. The somewhat erratic appearance of the genes within these species and their presence on a plasmid in some enteric bacteria suggests horizontal gene transmission.

Within the *Shigella* species, the genes for aerobactin synthesis are chromosomal and are closely linked. Hybridization studies indicate the genes are contained on a single *Eco*RI restriction endonuclease fragment of *S. flexneri*, *S. boydii*, and *S. sonnei* DNA. These sequences show considerable homology with the ColV aerobactin sequences as they hybridize under stringent conditions. Although *S. sonnei* contains the genes for aerobactin synthesis as indicated by hybridization to the ColV aerobactin sequences, no detectable aerobactin was produced. Members of this species synthesize enterobactin and thus have the genetic information for two separate iron transport systems. The presence of a functional phenolate iron transport system may have permitted the accumulation of mutations in the aerobactin biosynthetic genes of *S. sonnei*. The genes encoding the aerobactin receptor and aerobactin utilization in *S. sonnei* have remained intact since *S. sonnei* is able to transport and utilize exogenous aerobactin. Although *S. sonnei* has been reported to synthesize a hydroxamate distinct from aerobactin and one strain of *S. boydii* was reported to make a phenolate siderophore (25), this was not observed in any of the strains we tested.

Unlike the other *Shigella* species, *S. dysenteriae* lacks the genes for the aerobactin iron transport system. Chemical and bioassays were negative for this compound, and sequences homologous to the ColV aerobactin genes were not detected. Since the aerobactin genes are found in all the other *Shigella* species, it is possible that the genes were present at one time in *S. dysenteriae* but were lost by *IS1*-mediated deletion in the strains we assayed.

Restriction enzyme digestion and hybridization have demonstrated the presence of the insertion sequence *IS1* flanking the ColV aerobactin genes. Studies by McDougall and Neilands (18) and Perez-Casal and Crosa (24) have also shown that the ColV aerobactin genes are flanked by *IS1*.

The large number of copies of *IS1* in the *S. flexneri* chromosome makes it difficult to determine by hybridization whether any of the *IS1* copies are closely linked to the aerobactin genes. However, the high frequency of deletion mutants, and the changes in hybridization to both aerobactin genes and *IS1* in SA255, suggests that one or more copies of *IS1* are in close proximity to these genes.

The presence of these insertion sequences could promote transposition as well as deletion of the aerobactin genes (5, 11). In addition, *IS1* could provide a site for recombination between DNA containing the aerobactin genes and other DNA elements which have *IS1* (11). Since the aerobactin genes have been associated with increased virulence in *E. coli* strains (29), the spread of these genes through transposition or recombination is of potential medical significance.

ACKNOWLEDGMENTS

We are indebted to J. B. Neilands, A. Bindereif, and S. McDougall for providing the cloned aerobactin genes and for helpful discussions and to S. McIntire for providing the lambda carrying *IS1*. We thank J. Stoeber High for expert technical assistance.

This study was supported by grant F-941 from the Robert A. Welch Foundation.

LITERATURE CITED

1. Arnow, L. E. 1937. Colorimetric determination of the components of 3,4-dihydroxyphenylalanine-tyrosine mixtures. *J. Biol. Chem.* 118:531-537.
2. Atkin, C. L., J. B. Neilands, and H. J. Phaff. 1970. Rhodotorulic acid from species of *Leucosporidium*, *Rhodospiridium*, *Rhodotorula*, *Sporidiobolus*, and *Sporobolomyces*, and a new alanine-containing ferrichrome from *Cryptococcus melibiosum*. *J. Bac-*

- teriol. 103:722-733.
3. Bindereif, A., V. Braun, and K. Hantke. 1982. The cloacin receptor of ColV-bearing *Escherichia coli* is part of the Fe^{3+} -aerobactin transport system. *J. Bacteriol.* 150:1472-1475.
 4. Bindereif, A., and J. B. Neilands. 1983. Cloning of the aerobactin-mediated iron assimilation system of plasmid ColV. *J. Bacteriol.* 153:1111-1113.
 5. Cornelis, G., and H. Saedler. 1980. Deletions and an inversion induced by a resident IS1 of the lactose transposon Tn951. *Mol. Gen. Genet.* 178:367-374.
 6. Csaky, T. Z. 1948. On the estimation of bound hydroxylamine in biological materials. *Acta Chem. Scand.* 2:450-454.
 7. Dempsey, W. B., and N. S. Willetts. 1976. Plasmid co-integrates of prophage lambda and R factor R100. *J. Bacteriol.* 126:166-176.
 8. Formal, S. B., P. Gemski, Jr., L. S. Baron, and E. H. Lebrech. 1970. Genetic transfer of *Shigella flexneri* antigens to *Escherichia coli* K-12. *Infect. Immun.* 1:279-287.
 9. Gibson, F., and D. I. Magrath. 1969. The isolation and characterization of a hydroxamic acid (aerobactin) formed by *Aerobacter aerogenes* 62-1. *Biochim. Biophys. Acta* 192:175-184.
 10. Hansen, J. B., and R. H. Olsen. 1978. Isolation of large bacterial plasmids and characterization of the P2 incompatibility group plasmids pMG1 and pMG5. *J. Bacteriol.* 135:227-238.
 11. Kleckner, N. 1981. Transposable elements in prokaryotes. *Annu. Rev. Genet.* 15:341-404.
 12. Laird, A. J., D. W. Ribbons, G. C. Woodrow, and I. G. Young. 1980. Bacteriophage Mu-mediated gene transposition and in vitro cloning of the enterochelin gene cluster of *Escherichia coli*. *Gene* 11:347-357.
 13. Laird, A. J., and I. G. Young. 1980. Tn5 mutagenesis of the enterochelin gene cluster of *Escherichia coli*. *Gene* 11:359-366.
 14. Luke, R. K. J., and F. Gibson. 1971. Location of three genes concerned with the conversion of 2,3-dihydroxybenzoate into enterochelin in *Escherichia coli* K-12. *J. Bacteriol.* 107:557-562.
 15. Maniatis, T., E. F. Fritsch, and J. Sambrook. 1982. Molecular cloning, a laboratory manual. Cold Spring Harbor Laboratory, Cold Spring Harbor, N.Y.
 16. Maniatis, T., A. Jeffrey, and D. G. Kleid. 1975. Nucleotide sequence of the rightward operator of phage λ . *Proc. Natl. Acad. Sci. U.S.A.* 72:1184-1188.
 17. Marmur, J. 1961. A procedure for the isolation of deoxyribonucleic acid from micro-organisms. *J. Mol. Biol.* 3:208-218.
 18. McDougall, S., and J. B. Neilands. 1984. Plasmid- and chromosome-coded aerobactin synthesis in enteric bacteria: insertion sequences flank operon in plasmid-mediated systems. *J. Bacteriol.* 159:300-305.
 19. Neilands, J. B. 1981. Iron absorption and transport in microorganisms. *Annu. Rev. Nutr.* 1:27-46.
 20. Nyman, K., K. Nakamura, H. Ohtsubo, and E. Ohtsubo. 1981. Distribution of the insertion sequence IS1 in gram-negative bacteria. *Nature (London)* 289:609-612.
 21. Ohtsubo, H., K. Nyman, W. Doroszkiewicz, and E. Ohtsubo. 1981. Multiple copies of iso-insertion sequences of IS1 in *Shigella dysenteriae*. *Nature (London)* 291:640-643.
 22. Payne, S. M. 1980. Synthesis and utilization of siderophores by *Shigella flexneri*. *J. Bacteriol.* 143:1420-1424.
 23. Payne, S. M., D. W. Niesel, S. S. Peixotto, and K. M. Lawlor. 1983. Expression of hydroxamate and phenolate siderophores by *Shigella flexneri*. *J. Bacteriol.* 155:949-955.
 24. Perez-Casal, J. F., and J. H. Crosa. 1984. Aerobactin iron uptake sequences in plasmid ColV-K30 are flanked by inverted IS1-like elements and replication regions. *J. Bacteriol.* 160:256-265.
 25. Perry, R. D., and C. L. San Clemente. 1979. Siderophore synthesis in *Klebsiella pneumoniae* and *Shigella sonnei* during iron deficiency. *J. Bacteriol.* 140:1129-1132.
 26. Pollack, J. R., B. N. Ames, and J. B. Neilands. 1970. Iron transport in *Salmonella typhimurium*: mutants blocked in the biosynthesis of enterobactin. *J. Bacteriol.* 104:635-639.
 27. Rogers, H. J. 1973. Iron-binding catechols and virulence in *Escherichia coli*. *Infect. Immun.* 7:445-456.
 28. Southern, E. M. 1975. Detection of specific sequences among DNA fragments separated by gel electrophoresis. *J. Mol. Biol.* 98:503-517.
 29. Warner, P. J., P. H. Williams, A. Bindereif, and J. B. Neilands. 1981. ColV plasmid-specified aerobactin synthesis by invasive strains of *Escherichia coli*. *Infect. Immun.* 33:540-545.